

Synergie optique/radar pour l'estimation de l'évapotranspiration et de sa partition en vue d'un produit "stress hydrique de la plante"

Gilles Boulet, Zoubair Rafi, Valérie Le Dantec, Pascal Fanise, Albert Olioso, Salah Er-Raki et Olivier Merlin



Context and objectives

- Need to monitor **plant water use** as well as **plant water stress** for many applications :
 - Optimization of irrigation scheduling, water deficit agriculture...
 - Assess vulnerability of rainfed agrosystems to drought
- Dual Source Energy Balance models compute both evaporation E and transpiration T from TIR data, as well as potential E and T
- But they derive **2 unknowns** (E+T) out of the **sole** surface temperature (T_{surf})
 - How robust/reliable are total and component flux (T, E) and water stress retrievals for all situations in view of operational TIR products ?
 - Can we constraint ET and E/T with additional information such as surface soil moisture (from S1 data for example) to get rid of underdeterminancies ?
- Context = High resolution (50-100m) frequent revisit (2/3 days) TIR satellite mission TRISHNA (CNES/ISRO) as well as Copernicus LSTM

Soil Plant Atmosphere Remote Sensing Evapotranspiration (SPARSE) model

Same rationale as the TSEB model > How one gets **2 unknowns** (E i.e.LE_s, T i.e. LE_v) from **1 data source** (T_{surf}) ?



Consequence:

Moderately dry topsoil with a moderately stressed vegetation interpreted as a fully transpiring vegetation and a very dry soil



Saadi et al., HESS 2018

E (i.e. LE_s) forced by surface soil moisture



(Merlin et al., 2011)

Insitu capacitive probe for now Maybe S1 later on ?

Rainfed wheat in Tunisia (Kairouan)



E (i.e. LE_s) forced by surface soil moisture



deficit irrigation wheat site in Morocco ET from T_{surf} only or T_{surf} & θ_{surf}

Total water stress 1-LE/LE_{pot}



Total water stress 1-LE/LE_{pot}





Plant water stress 1-T/T_{pot}



Take home messages

- Water stress <u>of the whole surface</u> is fairly well retrieved by the uncalibrated model with a **0.2 uncertainty**, but many outliers challenge the robustness of any TIR ETR product (e.g. in a data assimilation or temporal interpolation perspective)
- Constraint by **surface soil moisture** improves total ET , T/ET as well as total and plant water stress for late mid-growth stage
- What about other indices linked to the **photosynthetic activity** (PRI, fluorescence...) or SWIR for midseason ?
- Need to find a way to evaluate **plant water stress** with additional measurements
- Extension to other sites with sapflow meas., as well as L8/S2/S1 data

